ELSEVIER

Toxicology 134 (1999) 217–218

TOXICOLOGY

Contents of volume 134

Methods for biological monitoring of propylene oxide exposure in Fischer 344 rats
Steady state concentrations in blood and hemoglogin adducts are used to arrive at an estimate of exposure level
S. Osterman-Golkar, H. Licea Pérez, G.A. Csanády, W. Kessler, J.G. Filser (Sweden, Germany) 1–8

Are circulating cytokines interleukin-6 and tumor necrosis factor α involved in chlorpyrifos-induced fever?
The role of IL-6 remains uncertain
C.J. Gordon, P.J. Rowsey (USA) 9–17

Lead inhibits *meso*-2,3-dimercaptosuccinic acid induced calcium transients in cultured rhesus monkey kidney cells
Lead alone does not increase intracellular calcium levels, but may block transient increases
P.L. Pokorski, M.J. McCabe, Jr., J.G. Pounds (USA) 19–26

Effect of lowered temperature on the toxicity of sulphur mustard in vitro and in vivo
Lower temperatures are protective and increase the therapeutic window
T.W. Sawyer, D. Risk (Canada) 27–37

Increased radiation-induced apoptosis in mouse thymus in the absence of metallothionein
Metallothionein is radioprotective
D. Xi Deng, L. Cai, S. Chakrabarti, M.G. Cherian (Canada) 39–49

Adriamycin-induced changes of creatine kinase activity in vivo and in cardiomyocyte culture
Inactivation of creatinbe kinase may be a primary event in adriamycin cardiotoxity
S.M. DeAtley, M.Y. Aksenov, M.V. Aksenova, B. Jordan, J.M. Carney, D.A. Butterfield (USA) 51–62

Elevated immunoglobulin E (IgE) levels in children with exposure to environmental lead
Lead exposure in children is suggested to increase IgE production
P.M. Lutz, T.J. Wilson, J. Ireland, A.L. Jones, J.S. Gorman, N.L. Gale, J.C. Johnson, J.E. Hewett (USA). 63–78

Persistent suppression of delayed-type hypersensitivity in adult F344 rats after perinatal exposure to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin
Perinatal exposure to TCDD induces a persistent depression of delayed-type response in adult rats
B.C. Gehrs, R.J. Smialowicz (USA) 79–88

Announcement 89

Instructions to authors 91

Review

Factors influencing the carcinogenicity of food chemicals
M.M. Manson, D.J. Benford (UK) 93–108

Doxorubicin-resistant LoVo adenocarcinoma cells display resistance to apoptosis induction by some but not all inhibitors of ser/thr phosphatases 1 and 2A
Development of multidrug resistane protects LoVoDx cells against okudaic but not cantharidic acid
S. Sieder, E. Richter, K. Becker, R. Heins, H.J. Steinfelder (Germany) 109–115

The antioxidant EPC-K1 attenuates NO-induced mitochondrial dysfunction, lipid peroxidation and apoptosis in cerebellar granule cells
Treatment with the antioxidant reverse the changes in cellular functions to a varying degree
T. Wei, C. Chen, J. Hou, B. Zhao, W. Xin, A. Mori (China, Japan) 117–126

Evidence for the presence of a microsomal NADH-dependent enzyme system that can bioactivate aromatic amines in the liver of rats and mice
A NADH-dependent enzyme is indirectly distinguished from the 'regular' P450 in a few in vivo and in vitro parameters
T. Marczylo, C. Ioannides (UK) 127–141
Environmental lead exposure and activity of δ-aminolevulinic acid dehydratase (ALA-D) in maternal and cord blood
Low exposures in 240 urban mothers show effects on this diagnostic enzyme
D. Campagna, G. Huel, F. Girard, J. Sahuquillo, P. Blot (France) 143–152
Binding of the potent allergen hexahydrophthalic anhydride in the mucosa of the upper respiratory and alimentary tract following single inhalation exposures in guinea pigs and rats
Inhalation of labeled HHPA leads to presence of radioactivity in various tissues
C.H. Lindh, B.A.G. Jönsson, G. Johannesson, X.-D. Zhang, H. Welinder, E.B. Brittebo (Sweden) 153–168
Success of pyridostigmine, physostigmine, eptastigmine and phosphotriesterase treatments in acute sarin intoxication
The esterase inhibitor plus the organophosphorus pesticide hydrolase protect against sarin intoxication
K. Tuovinen, E. Kaliste-Korhonen, F.M. Raushel, O. Hänninen (Finland, USA) 169–178
Dose–response hepatotoxicity of the peroxisome proliferator, perfluorodecanoic acid and the relationship to phospholipid metabolism in rats
Hepatic metabolic response is a characteristic response to the agent and not simply the result of cytotoxicity
M. Adinehzadeh, N.V. Reo, B.M. Jarnot, C.A. Taylor, D.R. Mattie (USA) 179–195
Cytokines, growth factors, and oxidative stress in HepG2 cells treated with ethanol, acetaldehyde, and LPS
The response pattern differs according to the toxic agent and times of exposure
Ma.C. Gutiérrez-Ruiz, S.C. Quiroz, V. Souza, L. Bucio, E. Hernández, I.P. Olivares, L. Llorente, F. Vargas-Voráckova, D. Kershenobich (Mexico) 197–207

Author index of volume 134 211

Subject index of volume 133 213–215

Contents of volume 133 217–218